Food Allergy Action Plan

Student's Name:D.O.B:	Tea	cher:		Place Child's	
ALLERGY TO:			Picture Here		
Asthmatic Yes* No *Higher risk fo	or severe reaction			Here	
♦ STEP 1: TREATMENT ♦					
Symptoms:			d Medication ²	determined	
• If a food allergen has been ingested, but <i>no symptoms</i> :		□ EpiPen	Antihistamine by physician authorizing treatment		
 Mouth Itching, tingling, or swelling of lips, tongue, mouth 		□ EpiPen	☐ Antihistamine ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐ ☐		
• Skin Hives, itchy rash, swelling of the face or extremities		☐ EpiPen	☐ Antihistamine		
 Gut Nausea, abdominal cramps, vomiting, diarrhea 		☐ EpiPen	☐ Antihistamine		
 Throat + Tightening of throat, hoarseness, hacking cough 		☐ EpiPen	☐ Antihistamine		
 Lung t Shortness of breath, repetitive coughing, wheezing 		☐ EpiPen	☐ Antihistamine		
■ Heart † Thready pulse, low blood pressure, fainting, pale, blueness		☐ EpiPen	☐ Antihistamine		
Other t		☐ EpiPen	☐ Antihistamine		
■ If reaction is progressing (several of the above areas affected), give		☐ EpiPen	☐ Antihistamine		
The severity of symptoms can quickly change. † Potentially list	fe-threatening.				
DOSAGE Epinephrine: inject intramuscularly (circle one) EpiPen EpiPen Jr. (see reverse side for instructions)					
Antihistamine: give					
Other: give					
medication/dose/route					
◆ STEP 2: EMERGENCY CALLS ◆					
1. Call 911 (or Rescue Squad:). State that an allergic reaction has been treated, and additional epinephrine may be needed)					
2. Dr at					
3. Emergency contacts: Name/Relationship Phone Number(s)					
a 1.)			2.)		
b 1.)		2.)			
c 1.)		2.)			
EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE CHILD TO MEDICAL FACILITY!					
Parent/Guardian Signature		Date			

Date_

Doctor's Signature_

(Required)

TRAINED STAFF MEMBERS				
1	Room			
2	Room			
3	Room			

EPIPEN® AND EPIPEN® JR. DIRECTIONS

Pull off gray activation cap.



Hold black tip near outer thigh (always apply to thigh).



- Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. Remove the EpiPen® unit and massage the injection area for 10 seconds.
- Once EpiPen® is used, call the Rescue Squad. State additional epinephrine may be needed. Take the used unit with you to the Emergency Room. Plan to stay for observationat the Emergency Room for at least 4 hours.

For children with multiple food allergies, consider providing separate Action Plans for different foods.



^{**}Medication checklist adapted from the Authorization of Emergency Treatment form developed by the Mount Sinai School of Medicine. Used with permission.